

در این مجموعه سوالات اولین المپیاد ریاضی به همراه پاسخ نهایی آن را مشاهده می کنید.

Problem A1) Prove that $(21n+4)/(14n+3)$ is irreducible for every natural number n .

پاسخ :

$$3(14n+3) - 2(21n+4) = 1.$$

Problem A2) For what real values of x is $\sqrt{x + \sqrt{2x-1}} + \sqrt{x - \sqrt{2x-1}} = A$, given (a) $A = \sqrt{2}$, (b) $A = 1$, (c) $A = 2$, where only non-negative real numbers are allowed in square roots and the root always denotes the non-negative root?

پاسخ :

(a) any x in the interval $[1/2, 1]$; (b) no solutions; (c) $x=3/2$.

حل :

Note that we require $x \geq 1/2$ to avoid a negative sign under the inner square roots. Since $(x-1)^2 \geq 0$, we have $x \geq \sqrt{2x-1}$, so there is no difficulty with $\sqrt{x - \sqrt{2x-1}}$, provided that $x \geq 1/2$.

Squaring gives $2x + 2\sqrt{(x^2-2x+1)} = A^2$. Note that the square root is $|x-1|$, not simply $(x-1)$. So we get finally $2x + 2|x-1| = A^2$. It is now easy to see that we get the solutions above.

Problem A3) Let a, b, c be real numbers. Given the equation for $\cos x$:

$$a \cos^2 x + b \cos x + c = 0,$$

form a quadratic equation in $\cos 2x$ whose roots are the same values of x . Compare the equations in $\cos x$ and $\cos 2x$ for $a=4, b=2, c=-1$.

حل :


You need that $\cos 2x = 2 \cos^2 x - 1$. Some easy manipulation then gives:

$$a^2 \cos^2 2x + (2a^2 + 4ac - 2b^2) \cos 2x + (4c^2 + 4ac - 2b^2 + a^2) = 0.$$

The equations are the same for the values of a, b, c given. The angles are $2\pi/5$ (or $8\pi/5$) and $4\pi/5$ (or $6\pi/5$).

Problem B1) Given the length $|AC|$, construct a triangle ABC with $\angle ABC = 90^\circ$, and the median BM satisfying $BM^2 = AB \cdot BC$.

: حل




Area = $AB \cdot BC / 2$ (because $\angle ABC = 90^\circ = BM^2 / 2$ (required) = $AC^2 / 8$ (because $BM = AM = MC$), so B lies a distance $AC/4$ from AC . Take B as the intersection of a circle diameter AC with a line parallel to AC distance $AC/4$.

Problem B2) An arbitrary point M is taken in the interior of the segment AB . Squares $AMCD$ and $MBEF$ are constructed on the same side of AB . The circles circumscribed about these squares, with centers P and Q , intersect at M and N .

- prove that AF and BC intersect at N ;
- prove that the lines MN pass through a fixed point S (independent of M);
- find the locus of the midpoints of the segments PQ as M varies.

: حل



(a) $\angle ANM = \angle ACM = 45^\circ$. But $\angle FNM = \angle FEM = 45^\circ$, so A, F, N are collinear. Similarly, $\angle BNM = \angle BEM = 45^\circ$ and $\angle CNM = 180^\circ - \angle CAM = 135^\circ$, so B, N, C are collinear.

(b) Since $\angle ANM = \angle BNM = 45^\circ$, $\angle ANB = 90^\circ$, so N lies on the semicircle diameter AB . Let NM meet the circle diameter AB again at S . $\angle ANS = \angle BNS$ implies $AS = BS$ and hence S is a fixed point.

(c) Clearly the distance of the midpoint of PQ from AB is $AB/4$. Since it varies continuously with M, it must be the interval between the two extreme positions, so the locus is a segment length $AB/2$ centered over AB.

Problem B3) The planes P and Q are not parallel. The point A lies in P but not Q, and the point C lies in Q but not P. Construct points B in P and D in Q such that the quadrilateral ABCD satisfies the following conditions: (1) it lies in a plane, (2) the vertices are in the order A, B, C, D, (3) it is an isosceles trapezoid with AB parallel to CD (meaning that $AD = BC$, but AD is not parallel to BC unless it is a square), and (4) a circle can be inscribed in ABCD touching the sides.

بدون پاسخ

www.SignalS.ir

E-mail : SignalS_ir@yahoo.com